

Suriname

Population 2017

563 402

World Bank Income Level 2017 Revision

Upper Middle Income

Estimated Cumulative Cost of Obesity 2017-2025 (US\$M) **505.5**





Childhood obesity* exceeds 10% in 7 of 11 CARICOM countries with data

Obesity in childhood is linked to high blood pressure, type II diabetes, and low self esteem

Childhood obesity tracks into adulthood increasing the likelihood of developing NCDs

*Based on the WHO definition of childhood obesity

Caribbean Policy En	vironmont

Fiscal Policies	3 /20	SSB Taxes in Barbados, Bermuda & Dominica
School Policies banning SSBs in schools	2 /20	Bermuda and Trinidad & Tobago

Marketing Regulation/
Mandatory Front of Pack 0/20
Nutrition Labeling

Nutrition Guidelines & Policies/ Healthy Schools

Multiple countries

Policies

CARICOM Childhood Obesity Prevention action frameworks:

- 1. HCC Civil Society Action Plan 2017-2021: Preventing Childhood Obesity in the Caribbean
- CARPHA Plan of Action for promoting healthy weights in the Caribbean: Prevention and control of childhood obesity 2014-2019
- 3. CARPHA 6-point policy package
- 4. PAHO Plan of Action for the prevention of obesity in children and adolescents 2014-2019

OBESITY AND OVERWEIGHT AMONG CHILDREN



WHO 2016*

Childhood levels of

OVERWEIGHT or **OBESE**

AGE	BOTH SEXES
5 to 9	34.4%
10 to 19	29.6%

31.1% BOTH SEXES AGE: 5 to 19



GSHS 2016**

Adolescent

OVERWEIGHT

AGE	MALE	FEMALE
13 to 15	28%	29.1%

BOTH SEXES 28.6%

Childhood levels of

OBESITY

AGE	BOTH SEXES
5 to 9	17.6%
10 to 19	12.3%

13.9% BOTH SEXES AGE: 5 to 19



Adolescent

OBESE

AGE	MALE	FEMALE
13 to 15	12.2%	11.1%

BOTH SEXES 11.6%

RISK FACTOR STATUS

Insufficient activity*** 2010

20.0

AGE	MALE	FEMALE
11 to 17	77.9%	83.9%

80.9%BOTH SEXES
AGE: 11 to 17





BOTH SEXES

\$\$	Attendance at physical education (PE) class on >= 3 days/week during the school year	33.1%
<u> </u>	Students who spent three or more hours per day during a typical or usual day doing sitting activities	44.5%
	Soft drink consumption STUDENTS (13-15 years) who usually drank carbonated soft drinks one or more times per day during the past 30 days (%)	79.1%
÷	Infant breastfeeding	2.8%

^{***}Percent of defined population attaining less than 60 minutes of moderate- to vigorous-intensity physical activity daily.

ADULT OVERWEIGHT AND OBESITY PREVALENCE

2016

Age-standardized prevalence of **OVERWEIGHT** and **OBESITY**

MALE FEMALE **53.4% 64.2%**

Age-standardized prevalence of **OBESITY**

MALE	FEMALE
18.9%	33.7%

58.9% BOTH SEXES

26.4% BOTH SEXES







Fresh fruit and vegetable intake is low in both Caribbean men and women, with less than 15% in most countries eating the recommended five or more portions a day.

References obesity and overweight among children:

*WHO. The values provided by the World Health Organization's Global Health Observatory Data repository are best estimates using methodologies that allows comparability across countries and time. The estimates are updated as more recent or revised data become available, or when there are changes to the methodology being used. Therefore, they are not always the same as official national estimates.

**GSHS. These estimates are based on cross-sectional surveys carried out in the countries and span the years 2007 to 2017. Given the time span of the data collection, the numbers quoted for some of the earlier years may not accurately reflect the current situation and comparisons between countries based on these figures is challenging.

References:

- 1. WHO: http://www.worldometers.info
- 2. **United Nations**, World Population Prospects, 2017 revision
- 3. Global, regional and national prevalence of overweight and obesity in children and adults 1980-2013: A systematic analysis Lancet. 2014 Aug 30; 384(9945): 766-781.
- 4. Global Health Observatory data repository
- 5. Global school based student health survey (Results for students aged 13-15 years)
- 6. World Health Organization NCD Progress Monitor, 2017.
- 7. **World Obesity Federation:** http://www.obesityday.worldobesity.org/ourdata2017 http://docs.wixstatic.com/ugd/6599c5_2a8fa0968efd4854b74f399abb3078c0.pdf?index=true
- 8. El estado de las guías alimentarias basadas en alimentos en América Latina y el Caribe: 21 años después de la Conferencia Internacional sobre Nutrición
- 9. **Risk factor surveys of individual CARICOM countries**. STEPS surveys were used for all countries except Barbados where HOTN was used and Jamaica where Jamaica Health and Lifestyle survey 2008 was used.

CHILDHOOD OBESITY PREVENTION SCORECARD (COPS)

WHO Recommends a comprehensive approach to prevention including implementation of key policies including: Taxation of SSBs; Regulation of Marketing to Children; Mandatory Front of Package Nutritional Labelling; and School Policies. The HCC CSAP Childhood Obesity Prevention Scorecard (COPS) has 14 indicators to measure a country's response to this emerging crisis.



CSAP: Civil Society Action Plan | COP: Childhood Obesity Prevention | SSBs: Sugar Sweetened Beverages | EDNP: Energy Dense Nutrient Poor

PRIORITIES AREAS



Trade and fiscal policies



literacy



Early childhood nutrition



Marketing of healthy and unhealthy foods and beverages to children



School- and community-based interventions



Resource mobilisation



Strategic planning, monitoring, and evaluation

Link to the real time COPS webpage on the HCC website at:

www.healthycaribbean.org/cops/
You will find resources such as policy documents for countries reporting progress